

Hafaliadau Cydamserol

1.

Datrysych yr hafaliadau cydamserol canlynol gan ddefnyddio dull algebraidd (nid graffigol). Rhaid i chi ddangos eich holl waith cyfrifo. [4]

$$4x - 3y = 11$$

$$6x - 2y = 9$$

2.

Datrysych yr hafaliadau cydamserol canlynol gan ddefnyddio dull algebraidd.
Rhaid i chi ddangos eich holl waith cyfrifo.

[4]

$$3x + 5y = 6.5$$

$$2x - 2y = -9$$

[4]

[illegible]

Mae Levi yn berchen ar far byrbryd.
Mae'r brechdanau i gyd â'r un pris ac mae'r diodydd i gyd â'r un pris.



Mae Levi'n ysgrifennu'r hafaliadau cydamserol canlynol:


$$\begin{aligned} 3x + 2y &= 720 \\ 2x + 5y &= 810. \end{aligned}$$

- Mae *y* yn cynrychioli

- [illegible]

5.

Mae dimensiynau a pherimedrau dau betryal, G a H, yn cael eu dangos isod.



Nid yw'r diagramau wedi'u lluniadu wrth raddfa

Mae Freya wedi ysgrifennu hafaliad ar gyfer petryal G:

$$10a + 16b = 96$$

- (a) Ysgrifennwch hafaliad ar gyfer petryal H. [1]

.....

- (b) Defnyddiwch dull algebraidd i gyfrifo a a b .
Trwy hynny, cyfrifwch ddimensiynau petryal H . [5]

This image shows a full page of white paper with horizontal dashed lines, typical of primary school writing paper. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Hyd petryal H yw cm

Lled petryal H yw cm

6.

Mae'r teulu Davies eisiau prynu dodrefn gardd. Mae eu canolfan arddio leol yn stocio'r brand arbennig yr hoffen nhw ei gael.

Pris 1 fainc ardd a 4 cadair yw £310.

Pris 2 fainc ardd a 3 cadair yw £345.

Mae gan y teulu Davies £450 i'w wario.

Oes ganddyn nhw ddigon o arian i brynu 2 fainc ardd a 5 cadair?

Rhaid i chi ddangos eich holl waith cyfrifo.

[5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Aeth Iona a Dafydd i siopa.
Prynodd y ddau ohonyn nhw boteli o'r un cordial cyrens duon ac o'r un dŵr soda.
Prynodd Iona 2 fotel o gordial cyrens duon a 3 potel o ddŵr soda.
Prynodd Dafydd 4 potel o gordial cyrens duon ac 1 botel o ddŵr soda.
Gwariodd Iona £2.04 a gwariodd Dafydd £2.48.
Defnyddiwyd ddull algebraidd i gyfrifo cost potel o gordial cyrens duon a chost potel o ddŵr soda.

[5]

[illegible]

Cost potel o ddŵr soda yw

8.

Mae Tomos, Gwen a Jen yn ymweld â'r un stondin ffrwythau.

Mae Tomos yn prynu 4kg mafon a 5kg cyrens duon am gyfanswm cost o £38.

Mae Gwen yn prynu 6kg mafon a 3kg cyrens duon am gyfanswm cost o £39.

Defnyddiwch ddull algebraidd i ddarganfod faint yw'r cyfanswm mae Jen yn ei dalu am 5kg mafon a 7kg cyrens duon.

[6]

9.

Gwyddonydd yw David.

O ganlyniad i arbrawf, mae e'n darganfod cysylltiadau rhwng dau newidyn f a g .
Er mwyn ysgrifennu'r casgliad mae angen iddo ddarganfod gwerthoedd f a g .

Dyma'r cysylltiadau mae David yn eu darganfod:

- Mae dwywaith f wedi'i adio at dair gwaith g yn rhoi cyfanswm o 5
- Mae tair gwaith f wedi'i adio at bedair gwaith g yn rhoi cyfanswm o 4

Lluniwch bâr o hafaliadau cydamserol a datryswch nhw i ddarganfod gwerthoedd f a g .

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

[5]

Defnyddiwch ddull algebraidd i gyfrifo gwerthoedd a a b .

[6]

[illegible]

Cynllun Marcio

1.

Method to find the first variable	M1	Allow one slip (but not in the equated variable).
Correct first variable	A1	
Method to find the second variable	m1	
Correct second variable	A1	
	4	$x = 0.5, y = -3$

2.

Unit 2 GCSE Maths June 2015 Higher Tier	T I C K	M A R K	Comment
10. Method to find the first variable	✓	M1	Allow one slip (but not in the equated
Correct first variable	✓	A1	variable)
Method to find the second variable	✓	m1	FT 'their first variable'
Correct second variable	✓	A1	$x = -2, y = 2.5$

3.

Method to solve, e.g. equate variable with attempt to +/- as appropriate	M1	Allow 1 slip in non equated variable
First variable correct	A1	
Method to find the second variable	m1	
Second variable correct	A1	
	4	a = 4.2 b = 6

4.

6(a) (... the <u>cost</u> of a sandwich in <u>pence</u> AND (... the) <u>cost</u> of a drink in <u>pence</u>	B2	Do not accept 'sandwiches' and 'drinks' need 'cost'('price') and unit 'pence' B1 if correct (cost) but 'in pence' omitted, or for either statement correct, or for ' <i>sandwiches in pence and drinks in pence</i> ' B0 for 'sandwiches' and 'drinks'
(b) Method, e.g. equal coefficients	M1	Allow 1 slip in non-equalised variable
Correct first variable	A1	$x = 180$ or $y = 90$
Method to find 2 nd variable, e.g. substitution	m1	
Correct second variable	A1	

5.

(a) $8a + 8b = 69.6$ or $a + b = 8.7$ or equivalent	B1	Accept unsimplified equations Mark final answer unless no other marks awarded in (b)
(b) Method to equate coefficients (allow 1 slip) with attempt to subtract First variable Method to find second variable Second variable (Length of rectangle H is) 23.1 (cm) AND (Width of rectangle H is) 11.7 (cm)	M1 A1 m1 A1 B1	OR alternative algebraic method (not trial & improvement), e.g. $a = 8.7 - b$ or $b = 8.7 - a$ substituted into $10a + 16b = 96$ FT for their equations in a similar format $a = 7.2$ $b = 1.5$ Accept length and width in either order FT provided both answers are positive and only provided M1, m1 awarded ($3a + b$ and $a + 3b$ correctly evaluated) <i>(FT use of $4a + 4b = 69.6$ leads to $a = 30.4$ and $b = -13$)</i>

6.

10. $b + 4c = 310$, $2b + 3c = 345$ Method to find first variable Correct first variable Correct second variable (£)455 AND 'No'	B1 M1 A1 A1 B1 5	Strategy of forming a pair of equations. (Do not penalise for not defining variables.) FT for equations of equivalent difficulty. $b = (£)90$, $c = (£)55$ OR $(£)345 + 2 \times (£)55$. FT 'their first variable' for second A1. FT 'their derived b and c'. An unsupported answer gets 0 marks.
---	---	--

7.

11. Equations $2b + 3s = 2(\cdot)04$ and $4b + (1)s = 2(\cdot)48$ Method, equating coefficients or alternative First variable correct Method to find second variable Second variable correct	S1 M1 A1 m1 A1	Accept other variables Allow 1 slip, but not in equated coefficients FT their equations in 2 variables provided at least 1 equation is correct Blackcurrant 54(p) or (£)0.54 Soda water 32(p) or (£)0.32 <i>Do not accept trial & improvement, maximum mark S1</i>
--	--------------------------------	---

8.

Linear GCSE Mathematics Higher Tier November 2015 Paper 1		FINAL MARK SCHEME Comments
12. $4r(\text{aspberries}) + 5b(\text{blackcurrants}) = (£)38$ $6r(\text{aspberries}) + 3b(\text{blackcurrants}) = (£)39$ Method to eliminate variable, e.g. equal coefficients First variable Substitute to find second variable Second variable Jen pays (£)50.5(0)	S1 M1 A1 m1 A1 B1 6	For both equations, in symbols or words, however '+' and '=' must be shown as symbols Do not accept $4\text{kg} + 5\text{kg} = 38$ with $6\text{kg} + 3\text{kg} = 39$ <i>FT provided at least 1 equation correct, but 1 slip in the other equation</i> Allow 1 error in one term, not one with equal coefficients raspberries $r = (£)4.5(0)$ or blackcurrants $b = (£)4$ FT their first variable FT for m1 not A1 if 'their first variable is negative' FT provided M1 and m1 awarded, for correct evaluation of 'their $5r + 7b$ '

9.

8. $2f + 3g = 5$ and $3f + 4g = 4$	B1	FT provided at least 1 correct equation and solving is of equivalent difficulty
Method, e.g. equal coefficients	M1	Allow 1 slip in non-equalised variable
Correct first variable	A1	$f = -8$ or $g = 7$
Method to find 2 nd variable, e.g. substitution	m1	
Correct second variable	A1	
	5	

10.

$10a + 15b = 180$	B1	Or equivalent
$4a + 105b = 270$ or $4a + 105b + 90 = 360$	B1	Or equivalent
Method to solve, e.g. equate one variable and intention to subtract	M1	FT provided equivalent level of difficulty and at least B1 previously awarded
		Allow 1 slip in one of the non equated variables
One variable correct	A1	a = 15
Method to find the second variable	m1	b = 2
Second variable correct	A1	Unsupported answers are not accepted
	6	Trial & improvement is not accepted