

Surname Atebion
First name(s)

Centre Number

Candidate Number
2



GCE A LEVEL

1305U60-1



S24-1305U60-1

WEDNESDAY, 12 JUNE 2024 – AFTERNOON

FURTHER MATHEMATICS – A2 unit 6
FURTHER MECHANICS B

1 hour 45 minutes

For Examiner's use only		
Question	Maximum Mark	Mark Awarded
1	11	
2	16	
3	13	
4	13	
5	13	
6	14	
Total	80	

ADDITIONAL MATERIALS

In addition to this examination paper, you will need:

- a Formula Booklet;
- a calculator.

INSTRUCTIONS TO CANDIDATES

Use black ink or black ball-point pen. Do not use gel pen or correction fluid.

You may use a pencil for graphs and diagrams only.

Write your name, centre number and candidate number in the spaces at the top of this page.

Answer **all** questions.

Take g as 9.8 ms^{-2} .

Write your answers in the spaces provided in this booklet. If you run out of space, use the additional page(s) at the back of the booklet, taking care to number the question(s) correctly.

INFORMATION FOR CANDIDATES

The maximum mark for this paper is 80.

The number of marks is given in brackets at the end of each question or part-question.

Sufficient working must be shown to demonstrate the **mathematical** method employed.

Answers without working may not gain full credit.

Unless the degree of accuracy is stated in the question, answers should be rounded appropriately.

You are reminded of the necessity for good English and orderly presentation in your answers.




JUN241305U60101

Reminder: Sufficient working must be shown to demonstrate the **mathematical** method employed.

1. Two smooth spheres A and B are moving on a smooth horizontal plane when they collide obliquely. When the spheres collide, the line joining their centres is parallel to the vector $\mathbf{j}$, as shown in the diagram below.

Immediately before the collision, sphere A has velocity $(6\mathbf{i} - 3\mathbf{j}) \text{ ms}^{-1}$ and sphere B has velocity $(-4\mathbf{i} + 7\mathbf{j}) \text{ ms}^{-1}$. Sphere A has mass 6 kg and sphere B has mass 2 kg .



Immediately after the collision, sphere B has velocity $(-4\mathbf{i} - 5\mathbf{j}) \text{ ms}^{-1}$.

- (a) Find the velocity of A immediately after the collision.

[3]

Egwyddor Cadwraeth Momentum (ar hyd llinell y candoau)

$$6 \times -3 + 2 \times 7 = 6 \times ? + 2 \times -5$$

$$-18 + 14 = 6 \times ? - 10$$

$$\frac{6}{6} = ?$$

$$? = 1$$

Egwyddor Cadwraeth Momentum (perpendicular i linell y candoau)

$$6 \times 6 + 2 \times -4 = 6 \times ? + 2 \times -4$$

$$36 - 8 = 6 \times ? - 8$$

$$\frac{36}{6} = ?$$

$$? = 6$$

Cyflymder A yn dilyn y gwrthdrawriad yw

$$6\mathbf{i} + \mathbf{j}$$



- (b) Calculate the coefficient of restitution between A and B.

[3]

Deddf Adferiad Newton (ar hyd llinell y canolau)

$$V_B - V_A = -e(u_B - u_A)$$

$$-5 - 1 = -e(7 - -3)$$


$$-6 = -e(10)$$

$$\frac{-6}{-10} = e$$

$$e = 0.6$$

- (c) Find the angle through which the direction of motion of B is deflected as a result of the collision. Give your answer correct to the nearest degree.

[3]



$$\tan \alpha = \frac{7}{4}$$

$$\alpha = \tan^{-1}\left(\frac{7}{4}\right)$$

$$\alpha = 60.2551187^\circ$$

$$\tan \beta = \frac{5}{4}$$

$$\beta = \tan^{-1}\left(\frac{5}{4}\right)$$

$$\beta = 51.34019175^\circ$$

Ongl allwyo

$$= 60.2551187^\circ + 51.34019175^\circ$$

$$= 111.5953105^\circ$$

$$= 112^\circ \text{ i'r radd agosaf.}$$



- (d) After the collision, sphere B continues to move with velocity $(-4\mathbf{i} - 5\mathbf{j}) \text{ ms}^{-1}$ until it collides with another sphere C , which exerts an impulse of $(-20\mathbf{i} + 18\mathbf{j}) \text{ Ns}$ on B .

Find the velocity of B after the collision with C .

[2]

Ergyd = Newid meun momentum

$$-20\mathbf{i} + 18\mathbf{j} = \text{mas (cyflymder uedyn - cyflymder gynt)}$$

$$-20\mathbf{i} + 18\mathbf{j} = 2(a\mathbf{i} + b\mathbf{j} - (-4\mathbf{i} - 5\mathbf{j}))$$

$$-20\mathbf{i} + 18\mathbf{j} = 2(a\mathbf{i} + b\mathbf{j} + 4\mathbf{i} + 5\mathbf{j})$$

$$-10\mathbf{i} + 9\mathbf{j} = a\mathbf{i} + b\mathbf{j} + 4\mathbf{i} + 5\mathbf{j}$$

$$-14\mathbf{i} + 4\mathbf{j} = a\mathbf{i} + b\mathbf{j}$$

Cyflymder B yn dilyn y gwrthdrawiad efo C yw
 $-4\mathbf{i} + 4\mathbf{j}$.



BLANK PAGE

**PLEASE DO NOT WRITE
ON THIS PAGE**




2. An object, of mass 1.8 kg , is falling vertically downwards under gravity. During the motion, it experiences a variable resistance of $0.2v^2 \text{ N}$, where $v \text{ ms}^{-1}$ is the speed of the object at time t seconds.

(a) Show that v satisfies the differential equation

$$\frac{dv}{dt} = \frac{9g - v^2}{9}$$

[3]

$$0.2v^2 \text{ N}$$



$$1.8g \text{ N}$$

$F = ma$, yn fertigol, $\downarrow = +$ if:

$$1.8g - 0.2v^2 = 1.8a$$

$$g - \frac{1}{9}v^2 = a$$

$$\frac{9g}{9} - \frac{1}{9}v^2 = \frac{dv}{dt}$$

$$\frac{9g - v^2}{9} = \frac{dv}{dt}$$

$$\frac{dv}{dt} = \frac{9g - v^2}{9} \quad \checkmark$$



At time $t = 0$, the object passes a point A with a speed of $\sqrt{g}$ ms⁻¹. The object then hits the ground with a speed of 8 ms⁻¹.

(b) Calculate the time taken for the object to hit the ground.

[6]

$$\frac{dv}{dt} = \frac{9g - v^2}{9}$$

$$\frac{9}{9g - v^2} dv = dt$$

$$\int \frac{1}{9g - v^2} dv = \int \frac{1}{9} dt$$

$$\int \frac{1}{(3\sqrt{g})^2 - v^2} dv = \int \frac{1}{9} dt$$

$$\frac{1}{3\sqrt{g}} \tanh^{-1}\left(\frac{v}{3\sqrt{g}}\right) = \frac{1}{9} t + C$$

As yw $t=0$, $v = \sqrt{g}$, felly

$$\frac{1}{3\sqrt{g}} \tanh^{-1}\left(\frac{\sqrt{g}}{3\sqrt{g}}\right) = \frac{1}{9}(0) + C$$

$$\frac{1}{3\sqrt{g}} \tanh^{-1}\left(\frac{1}{3}\right) = C$$

$$\text{Felly } \frac{1}{3\sqrt{g}} \tanh^{-1}\left(\frac{v}{3\sqrt{g}}\right) = \frac{1}{9} t + \frac{1}{3\sqrt{g}} \tanh^{-1}\left(\frac{1}{3}\right)$$

$$\text{As } yw \ v=8, \frac{1}{3\sqrt{g}} \tanh^{-1}\left(\frac{8}{3\sqrt{g}}\right) = \frac{1}{9} t + \frac{1}{3\sqrt{g}} \tanh^{-1}\left(\frac{1}{3}\right)$$

$$\frac{3}{\sqrt{g}} \tanh^{-1}\left(\frac{8}{3\sqrt{g}}\right) - \frac{3}{\sqrt{g}} \tanh^{-1}\left(\frac{1}{3}\right) = t$$

$$t = 0.8780376185$$

$$t = \underline{\underline{0.885}} \text{ ; 2 le degol}$$



- (c) Given that the distance of the object from A at time t is x metres, form another differential equation to find an expression for x in terms of v . Hence, find the height of A above the ground. [7]

Diagram: A point A is shown with a vertical line extending downwards to a horizontal line representing the ground. The distance from A to the ground is labeled h . The initial velocity is $\sqrt{g} \text{ ms}^{-1}$ and the final velocity is 8 ms^{-1} .

Rheol y gadwyn: $\frac{dv}{dt} = \frac{dx}{dt} \times \frac{dv}{dx}$
 $= v \times \frac{dv}{dx}$

Felly $v \frac{dv}{dx} = \frac{9g - v^2}{9}$

$\frac{9v}{9g - v^2} dv = dx$

$\int \frac{9v}{9g - v^2} dv = \int dx$

$-\frac{9}{2} \ln|9g - v^2| = x + C$

OS yw $x=0$, $v=\sqrt{g}$, felly
 $-\frac{9}{2} \ln|9g - (\sqrt{g})^2| = 0 + C$

$-\frac{9}{2} \ln|9g - g| = C$

$-\frac{9}{2} \ln(8g) = C$

Felly
 $-\frac{9}{2} \ln|9g - v^2| = x - \frac{9}{2} \ln(8g)$

$\frac{9}{2} \ln(8g) - \frac{9}{2} \ln|9g - v^2| = x$

$\frac{9}{2} \ln \left| \frac{8g}{9g - v^2} \right| = x$

OS yw $v = 8 \text{ ms}^{-1}$,
 $\frac{9}{2} \ln \left| \frac{8g}{9g - 8^2} \right| = x$

$x = \frac{9}{2} \ln \left| \frac{8 \times 9.8}{9 \times 9.8 - 64} \right|$

$x = \frac{9}{2} \ln \left(\frac{392}{121} \right)$

$x = 5.289620824$

$x = \underline{\underline{5.29 \text{ m}}}$ i 2 lc deeg




BLANK PAGE

**PLEASE DO NOT WRITE
ON THIS PAGE**



3. ACB is the diameter of a semi-circular lamina of radius $2a$ and centre C . Another semi-circular lamina, having AC as its diameter, is added to form a uniform lamina, as shown in the diagram below.



- (a) (i) Show that the distance of the centre of mass of the lamina from AB is $\frac{28}{15\pi}a$.

- (ii) Calculate the distance of the centre of mass of the lamina from a line drawn through A that is perpendicular to AB . [10]

Gwrthrych	Arwynebedd	Pellter craidd mäs o AB	Pellter craidd mäs o AD
$\frac{1}{2}$ cylch AB	$\frac{\pi(2a)^2}{2}$	$\frac{4(2a)}{3\pi}$	$2a$
$\frac{1}{2}$ cylch AC	$\frac{\pi a^2}{2}$	$-\frac{4a}{3\pi}$	a
Lamina	$\frac{\pi(2a)^2}{2} + \frac{\pi a^2}{2}$	$\bar{x}$	$\bar{y}$



(i) Yn cymryd momentau o amgylch AB:

$$\left(\frac{\pi(2a)^2}{2} + \frac{\pi a^2}{2} \right) \bar{x} = \frac{\pi(2a)^2}{2} \times \frac{4(2a)}{3\pi} + \frac{\pi a^2}{2} \times \frac{-4a}{3\pi}$$

$$\left(\frac{4a^2\pi}{2} + \frac{\pi a^2}{2} \right) \bar{x} = \frac{4a^2\cancel{\pi} \times 8a}{2 \times 3\cancel{\pi}} - \frac{4\cancel{\pi} a^3}{6\cancel{\pi}}$$

$$\frac{5a^2\pi}{2} \bar{x} = \frac{32a^3}{6} - \frac{4a^3}{6}$$

$$\frac{5a^2\pi}{2} \bar{x} = \frac{14a^3}{3}$$

$$\bar{x} = \frac{14\cancel{a^2} \times 2}{3 \times 5\cancel{a^2}\pi}$$

$$\bar{x} = \frac{28}{15\pi} a \quad \checkmark$$

(ii) Yn cymryd momentau o amgylch AD:

$$\frac{5a^2\pi}{2} \bar{y} = \frac{4a^2\pi}{2} \times \cancel{2}a + \frac{\pi a^2}{2} \times a$$

$$\frac{5a^2\pi}{2} \bar{y} = 4a^3\pi + \frac{1}{2}a^3\pi$$

$$5a^2\cancel{\pi} \bar{y} = 8a^3\cancel{\pi} + a^3\cancel{\pi}$$


$$5a^2 \bar{y} = 9a^3$$

$$\bar{y} = \frac{9a^3}{5a^2}$$


$$\bar{y} = \frac{9a}{5}$$



- (b) Suppose that the lamina is suspended in equilibrium by means of two vertical wires attached at A and B so that AB is horizontal. Find the fraction of the lamina's weight that is supported by the wire attached at B . [3]



Modelu fel rhodan anffurf:



Yn cymryd momentau o amgylch A :

$$\begin{aligned} \text{Momentau clocwedd} \\ = \text{Momentau gwrthglocwedd} \\ T_B \times 4a = mg \times \frac{9a}{5} \end{aligned}$$

$$T_B = \frac{9mg}{5 \times 4}$$

$$T_B = \frac{9mg}{20}$$

Y ffrasiun o'r pwysau sy'n cael ei gymal yn B yw $\frac{9}{20}$




BLANK PAGE

**PLEASE DO NOT WRITE
ON THIS PAGE**



4. The diagram below shows part of a game at a funfair that consists of a target moving along a straight horizontal line AB . The centre of the target may be modelled as a particle moving with Simple Harmonic Motion about centre O , where O is the midpoint of AB .



When the target is at a distance of 84 cm from O , its speed is 52 cm s^{-1} and the magnitude of its acceleration is 1344 cm s^{-2} .

- (a) Show that the period of the motion is $\frac{\pi}{2} \text{ s}$.

[3]

In defnyddio $\frac{d^2x}{dt^2} = -\omega^2 x$ gyda $\rightarrow = +$ if

$$1344 = -\omega^2(-84)$$

$$\frac{1344}{84} = \omega^2$$

$$16 = \omega^2$$

$$\omega = 4.$$

Nawr cyfnod = $\frac{2\pi}{\omega}$

$$= \frac{2\pi}{4}$$

$$= \frac{\pi}{2} \text{ s} \quad \checkmark$$



- (b) Determine the maximum speed of the target.

Examiner
only

[5]

Yn defnyddio $v^2 = w^2(a^2 - x^2)$

$$52^2 = 4^2(a^2 - (-84)^2)$$

$$2704 = 16(a^2 - 7056)$$

$$169 = a^2 - 7056$$

$$7225 = a^2$$

$$a = 85$$


Y buanedd macsimum yw a w (Car y canol)

$$= 85 \times 4$$

$$= 340 \text{ cms}^{-1}$$



- (c) During a game, players fire a ball at the target. A timer is started when the target is at A. Players must wait for the target to complete at least one full cycle before firing. Given that the target is hit when it is at a distance of 67 cm from O, calculate the two earliest possible times taken to hit the target. [5]




Yn defnyddio $x = a \sin(\omega t + \epsilon)$

gwelwn fod $\epsilon = \frac{3\pi}{2}$ gan fod y mudiant yn cychwyno'r chwith.

Felly $x = 85 \sin(4t + \frac{3\pi}{2})$

Os yw $x = -67$, yna $-67 = 85 \sin(4t + \frac{3\pi}{2})$
 $-\frac{67}{85} = \sin(4t + \frac{3\pi}{2})$



$$\sin^{-1}\left(-\frac{67}{85}\right) = 4t + \frac{3\pi}{2}$$

$$\dots, -0.9079, 4.0495, 5.3753, \dots = 4t + \frac{3\pi}{2}$$

$$\dots, -5.6203, -0.6629, 0.6629, \dots = 4t$$

$$\dots, -1.4051, -0.1657, 0.1657, \dots = t$$

Trwy gymesuredd y graff, y ddau amser cynharaf
 posib yw $\frac{\pi}{2} + 0.1657 = \underline{\underline{1.7365s}}$ (i 4 lle degol)

$$a \quad \frac{3\pi}{4} - 0.1657 = \underline{\underline{2.1905s}} \quad (\text{i 4 lle degol})$$




BLANK PAGE

**PLEASE DO NOT WRITE
ON THIS PAGE**



5. The diagram below shows a uniform rod AB of weight WN and length $2l$, with its lower end A resting on a rough horizontal floor. A light cable is attached to the other end B . The rod is in equilibrium when it is inclined at an angle of θ to the floor, where $0^\circ < \theta \leq 45^\circ$. The tension in the cable is TN acting at an angle of 2θ to the rod, as shown in the diagram.



- (a) (i) Show that $T = \frac{W}{4} \operatorname{cosec} \theta$. [4]
 (ii) Hence determine the normal reaction of the floor on the rod at A , giving your answer in terms of W . [4]

(c) Yn cymryd momentau o amgylch A :

Momentau cloedd = momentau gwrthglawedd

$$T \sin 2\theta \times 2l = W \cos \theta \times l$$


$$T(2 \sin \theta \cos \theta) \times 2 = W \cos \theta$$

$$4T \sin \theta = W$$

$$T = \frac{W}{4 \sin \theta}$$

$$T = \frac{W}{4} \operatorname{cosec} \theta$$

$$T = \frac{W}{4} \operatorname{cosec} \theta \quad \checkmark$$



Yn defnyddio grymoedd i fyny = grymoedd i lawr

$$R_A + T \sin \theta = W$$

$$R_A = W - T \sin \theta$$

$$R_A = W - \left(\frac{W \operatorname{cosec} \theta}{4} \right) \sin \theta$$

$$R_A = W - \left(\frac{W}{4 \sin \theta} \right) \sin \theta$$

$$R_A = W - \frac{W}{4}$$

$$R_A = \frac{3W}{4} \text{ N}$$



- (b) Given that the coefficient of friction between the floor and the rod is $\frac{\sqrt{3}}{3}$, calculate the minimum possible value for θ . [5]

Grymoedd i'r dde = grymoedd i'r chwith

$$T \cos \theta = F_{\text{friction}}$$

$$\left(\frac{W}{4} \operatorname{cosec} \theta \right) \cos \theta = \mu R_A \quad (\text{ar fin symud})$$

$$\left(\frac{W}{4 \sin \theta} \right) \cos \theta = \frac{\sqrt{3}}{3} \left(\frac{3W}{4} \right)$$

$$\frac{1}{4} \cot \theta = \frac{\sqrt{3}}{4}$$

$$\cot \theta = \sqrt{3}$$

$$\tan \theta = \frac{1}{\sqrt{3}}$$

$$\theta = \tan^{-1} \left(\frac{1}{\sqrt{3}} \right)$$


$$\underline{\underline{\theta = 30^\circ}}$$



6. The region R , shown in the diagram below, is bounded by the coordinate axes and the curve

$$y = \frac{a}{b} \sqrt{b^2 - x^2},$$

where a, b are constants.



The region R is rotated through 360° about the x -axis to form a uniform solid S .

The volume of S is $\frac{2}{3} \pi a^2 b$.

- (a) Use integration to show that the distance of the centre of mass of S from the y -axis is $\frac{3b}{8}$.

[5]

Mae'r solid yn unfurf, fel bodei ddwysedd ρ yn gyson.
(ρ yw'r mas fesul uned cyfaint.)

Gadewch i ni rannu'r solid yn dafellau yn baralel i'r echelin- y . Ystyriwch dafell PQ, sydd ar bellter x oddi wrth O a thrwydd δx .

Radiws y dafell yw $\frac{a}{b} \sqrt{b^2 - x^2}$.

Màs y dafell = cyfaint $\times$ dwysedd

$$= \pi r^2 \times \text{trwydd} \times \text{dwysedd}$$

$$= \pi \left(\frac{a}{b} \sqrt{b^2 - x^2} \right)^2 \times \delta x \times \rho$$

$$= \pi \frac{a^2}{b^2} (b^2 - x^2) \delta x \rho$$

Mae hwn yn gweithredu ar bellter x o O .



$$\begin{aligned}\text{Màs y solid} &= \text{cyfaint} \times \text{dwysedd} \\ &= \frac{2}{3} \pi a^2 b \rho\end{aligned}$$

Mae hwn yn gweithredu ar bellter $\bar{x}$ o 0.

Yn cymryd momentau o amgylch yr echelin-y:

$$\bar{x} = \frac{\sum (\text{Màs scribed} \times \text{Bellter scribed o 0})}{\text{Màs y solid}}$$


Màs y solid

$$(\text{Màs y solid}) \bar{x} = \sum (\text{Màs scribed} \times \text{Bellter scribed o 0})$$

$$\frac{2}{3} \pi a^2 b \rho \bar{x} = \sum \left(\pi \frac{a^2}{b^2} (b^2 - x^2) \delta x \rho \times x \right)$$

→ Wele tudalen 25

The diagram below shows a small tree growing in a pot. The uniform solid S described on the previous page may be used to model the part of the tree above the pot. This part of the tree has height h cm and base radius $\frac{h}{4}$ cm. The pot, including its contents, may be modelled as a solid cylinder of height 50 cm and radius 25 cm.



You may assume that the density of the pot, including its contents, is equal to 20 times the density of the part of the tree above the pot.



- (b) A gardener suggests that a tree is said to have outgrown its pot if the centre of mass, of both the tree and its pot, lies above the height of the pot. Determine the maximum value of h before the tree outgrows its pot. [8]

Gradewch i ρ gynrychioli dwysedd y goeden.

Siâp	Mäs	Pellter craidd mäs o'r gwaelod
Silindr	$(\pi r^2 h)(20\rho)$ $= \pi(25^2)(50)(20\rho)$ $= 625000\pi\rho$	25cm
Coeden	$\frac{2}{3}\pi a^2 b(\rho)$ $= \frac{2}{3}\pi\left(\frac{h}{4}\right)^2 h\rho$ $= \frac{2}{3}\pi\left(\frac{h^2}{16}\right)h\rho$ $= \frac{1}{24}\pi h^3\rho$	$50 + \frac{3b}{8}$ $= 50 + \frac{3h}{8}$ cm
Solid Cyfansawdd	$625000\pi\rho + \frac{1}{24}\pi h^3\rho$	$\bar{h}$

Yn cymryd momentau o amgylch y gwaelod:

$$(625000\pi\rho + \frac{1}{24}\pi h^3\rho)\bar{h} = 625000\pi\rho(25) + \frac{1}{24}\pi h^3\rho(50 + \frac{3h}{8})$$

$$(625000 + \frac{1}{24}h^3)\bar{h} = 15625000 + \frac{1}{24}h^3(50 + \frac{3h}{8})$$

$$(625000 + \frac{1}{24}h^3)\bar{h} = 15625000 + \frac{25}{12}h^3 + \frac{1}{64}h^4$$

Os yw $\bar{h} = 50$ yna

$$(625000 + \frac{1}{24}h^3)50 = 15625000 + \frac{25}{12}h^3 + \frac{1}{64}h^4$$

$$31250000 + \frac{25}{12}h^3 = 15625000 + \frac{25}{12}h^3 + \frac{1}{64}h^4$$

$$15625000 = \frac{1}{64}h^4$$

$$1,000,000,000 = h^4$$

$$h = \underline{\underline{177.83 \text{ cm}}} \text{ i 2 le degol}$$



- (c) Identify one possible limitation of the model used that could affect your answer to part (b).

[1]

Ni fyddai dwysedd y silindr yn unffurf, h.y. bydd
y gwriddiau'r goeden yn siapiau gwahanol wrth
fynd trwy'r pot.

END OF PAPER



Question
numberAdditional page, if required.
Write the question number(s) in the left-hand margin.Examiner
only

6a

Fel mae $\delta x \rightarrow 0$ mae hun yn rhoi

$$\frac{2}{3} \pi a^2 b \bar{x} = \int_0^b \frac{\pi a^2}{b^2} (b^2 - x^2) x dx$$

$$\frac{2}{3} \pi a^2 b \bar{x} = \pi \frac{a^2}{b^2} \int_0^b (b^2 - x^2) x dx$$

$$\frac{2}{3} b \bar{x} = \frac{1}{b^2} \int_0^b b^2 x - x^3 dx$$

$$\frac{2}{3} b \bar{x} = \frac{1}{b^2} \left[\frac{b^2 x^2}{2} - \frac{x^4}{4} \right]_0^b$$

$$\frac{2}{3} b \bar{x} = \frac{1}{b^2} \left[\left(\frac{b^2(b^2)}{2} - \frac{b^4}{4} \right) - \left(\frac{b^2(0^2)}{2} - \frac{0^4}{4} \right) \right]$$

$$\frac{2}{3} b \bar{x} = \frac{1}{b^2} \left[\frac{b^4}{2} - \frac{b^4}{4} - (0 - 0) \right]$$

$$\frac{2}{3} b \bar{x} = \frac{1}{b^2} \left(\frac{b^4}{4} \right)$$

$$\frac{2}{3} b \bar{x} = \frac{b^2}{4}$$

$$\bar{x} = \frac{3b^2}{4 \times 2b}$$

$$\bar{x} = \frac{3b}{8} \quad \checkmark$$



[illegible]

Examiner
only



BLANK PAGE

**PLEASE DO NOT WRITE
ON THIS PAGE**



BLANK PAGE

**PLEASE DO NOT WRITE
ON THIS PAGE**

