



GCE AS/A level

0977/01



S16-0977-01

MATHEMATICS – FP1
Further Pure Mathematics

A.M. FRIDAY, 24 June 2016

1 hour 30 minutes

ADDITIONAL MATERIALS

In addition to this examination paper, you will need:

- a 12 page answer book;
- a Formula Booklet;
- a calculator.

INSTRUCTIONS TO CANDIDATES

Use black ink or black ball-point pen.

Answer **all** questions.

Sufficient working must be shown to demonstrate the **mathematical** method employed.

INFORMATION FOR CANDIDATES

The number of marks is given in brackets at the end of each question or part-question.

You are reminded of the necessity for good English and orderly presentation in your answers.

1. Differentiate $\frac{x^2}{x+1}$ from first principles. [7]

2. The transformation T in the plane consists of an anticlockwise rotation through 90° about the origin followed by a translation in which the point (x, y) is transformed to the point $(x + 1, y + 2)$.

(a) Determine the 3×3 matrix which represents T . [4]

(b) Find the fixed point of T . [4]

3. Given that

$$S_n = \sum_{r=1}^n r^2(r+1),$$

obtain an expression for S_n in terms of n , giving your answer as a product of four linear factors. [6]

4. The complex numbers z_1, z_2 are given by

$$z_1 = -\sqrt{3} + i; \quad z_2 = 1 + i.$$

(a) Determine the modulus and the argument of each of z_1, z_2 , giving **exact** values of the moduli and giving the arguments in terms of π . [4]

(b) The complex number w is given by

$$w = \frac{z_1^2}{z_2}.$$

Using your results from (a), or otherwise, determine w in the form $a + ib$, giving a, b correct to two decimal places. [6]

5. The matrix \mathbf{M} is given by

$$\mathbf{M} = \begin{bmatrix} 2 & 5 & \lambda \\ 0 & \lambda & -1 \\ \lambda & 2 & 1 \end{bmatrix}.$$

(a) (i) Show that

$$\det \mathbf{M} = 4 - 3\lambda - \lambda^3.$$

(ii) Hence show that \mathbf{M} is singular when $\lambda = 1$ and is not singular for any other real values of λ .

(iii) Show that the following system of equations is consistent and find the general solution. [12]

$$\begin{bmatrix} 2 & 5 & 1 \\ 0 & 1 & -1 \\ 1 & 2 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \\ 1 \end{bmatrix}$$

(b) Suppose now that $\lambda = -1$. By first finding the adjugate matrix of \mathbf{M} , determine the inverse matrix \mathbf{M}^{-1} . [5]

6. Consider the cubic equation

$$ax^3 + bx^2 + cx + d = 0.$$

Given that the product of two of the roots is equal to 1, show that

$$d^2 - bd = a^2 - ac. \quad [6]$$

7. The sequence x_1, x_2, x_3, \dots is generated by the relationship

$$x_{n+1} = 2x_n - n + 1 \quad \text{where } x_1 = 3.$$

Use mathematical induction to prove that

$$x_n = 2^n + n$$

for all positive integers n . [6]

8. The function f is defined on the domain $\left(0, \frac{\pi}{2}\right)$ by

$$f(x) = x^{\sin x}.$$

(a) Obtain an expression for $f'(x)$. [4]

(b) Given that the graph of f has one stationary point, show that its x -coordinate lies between 0.35 and 0.36. [3]

TURN OVER.

9. The complex numbers z and w are represented, respectively, by points $P(x, y)$ and $Q(u, v)$ in Argand diagrams and

$$w = (z + 2i)^2.$$

- (a) Obtain expressions for u and v in terms of x and y . [4]
- (b) The point P moves along the line $y = x - 1$. Find the equation of the locus of Q . [4]

END OF PAPER