



**GCE AS/A level**

0981/01



S16-0981-01

**MATHEMATICS M2**

**Mechanics**

A.M. TUESDAY, 21 June 2016

1 hour 30 minutes

### **ADDITIONAL MATERIALS**

In addition to this examination paper, you will need:

- a 12 page answer book;
- a Formula Booklet;
- a calculator.

### **INSTRUCTIONS TO CANDIDATES**

Use black ink or black ball-point pen.

Answer **all** questions.

Take  $g$  as  $9.8 \text{ ms}^{-2}$ .

Sufficient working must be shown to demonstrate the **mathematical** method employed.

### **INFORMATION FOR CANDIDATES**

The number of marks is given in brackets at the end of each question or part-question.

You are reminded of the necessity for good English and orderly presentation in your answers.

1. A particle of mass 4 kg moves along the  $x$ -axis, starting, when  $t = 0$ , from the point where  $x = 3$ . At time  $t$  s, its velocity  $v \text{ ms}^{-1}$  is given by

$$v = 12t^2 - 7kt + 1,$$

where  $k$  is constant.

When  $t = 2$ , the displacement of the particle from the origin is 16 m.


- (a) Determine the value of  $k$ . [5]
- (b) Calculate the magnitude of the force acting on the particle when  $t = 5$ . [4]
2. A particle is projected from horizontal ground with speed  $24.5 \text{ ms}^{-1}$  in a direction inclined at an angle of  $30^\circ$  above the horizontal.
- (a) Calculate the horizontal range of the particle. [6]
- (b) Determine the maximum height reached by the particle. [3]
- (c) Write down the speed and the direction of motion of the particle as it hits the ground. [1]
3. At time  $t = 0$  s, the position vector of an object  $A$  is  $\mathbf{i}$  m and the position vector of another object  $B$  is  $3\mathbf{i}$  m. The constant velocity vector of  $A$  is  $2\mathbf{i} + 5\mathbf{j} - 4\mathbf{k} \text{ ms}^{-1}$  and the constant velocity vector of  $B$  is  $\mathbf{i} + 3\mathbf{j} - 5\mathbf{k} \text{ ms}^{-1}$ . Determine the value of  $t$  when  $A$  and  $B$  are closest together and find the least distance between  $A$  and  $B$ . [9]
4. By burning a charge, a cannon fires a cannon ball of mass 12 kg horizontally. As the cannon ball leaves the cannon, its speed is  $600 \text{ ms}^{-1}$ . The recoiling part of the cannon has a mass of 1600 kg.
- (a) Determine the speed of the recoiling part immediately after the cannon ball leaves the cannon. [3]
- (b) Find the energy created by the burning of the charge. State any assumption you have made in your solution. [4]
- (c) Calculate the constant force needed to bring the recoiling part to rest in 1.2 m. [2]
5. A particle is attached to one end of a light elastic string of natural length  $l$  m and modulus of elasticity  $\lambda$  N. The other end of the string is attached to the ceiling. The particle hangs in equilibrium. The length of the string is 0.95 m when the weight of the particle is 30 N, and 1.15 m when the weight of the particle is 70 N. Find the value of  $l$  and the value of  $\lambda$ . [6]

6. A particle moves on a horizontal plane such that its velocity vector  $v \text{ ms}^{-1}$  at time  $t$  s is given by

$$\mathbf{v} = 7 \sin 2t \mathbf{i} + 6 \cos 3t \mathbf{j}.$$

- (a) Find the acceleration vector of the particle at time  $t$  s. [2]
- (b) Given that when  $t = 0$ , the particle has position vector  $(0.5\mathbf{i} + 3\mathbf{j})$  m, find the position vector of the particle when  $t = \frac{\pi}{2}$ . [5]

7. The diagram below shows two points  $A$  and  $B$  on a mountain bike track.



The heights of  $A$  and  $B$  above ground level are 20 m and 22 m respectively. The length of the track between  $A$  and  $B$  is 16 m. The resistance to motion of a biker on the track may be modelled by a constant force of magnitude 50 N. The total mass of the biker and his bike is 70 kg. The speed of the biker at  $A$  is  $v \text{ ms}^{-1}$ . Find the minimum value of  $v$  if the biker is to reach  $B$  without pedalling. [7]

8. A rough circular plate rotates horizontally about a smooth fixed vertical axis through its centre  $O$ . A point  $A$  on the plate moves with constant speed  $v \text{ ms}^{-1}$ , where  $OA$  is 1.6 m. A particle of mass  $m$  kg lies on the point  $A$  on the plate. The coefficient of friction between the particle and the plate is 0.72. Given that the particle remains at the point  $A$ , find the greatest possible value of  $v$ . Hence write down the greatest possible value of the angular velocity of the plate. State clearly your units for the angular velocity. [7]
9. A smooth sphere, with centre  $O$  and radius 4 m, is fixed. A particle  $P$ , of mass  $m$ , resting on the sphere at its highest point, is given a horizontal speed of magnitude  $\sqrt{g} \text{ ms}^{-1}$ , where  $g$  is the magnitude of the acceleration due to gravity. At the instant the line  $OP$  makes an angle  $\theta$  with the upwards vertical, the speed of  $P$  is  $v \text{ ms}^{-1}$ .
- (a) Determine an expression for  $v^2$  in terms of  $g$  and  $\theta$  while  $P$  remains in contact with the sphere. [4]
- (b) Find, in terms of  $m$ ,  $g$  and  $\theta$ , the magnitude of the force exerted by the sphere on  $P$ . Hence calculate the value of  $\cos \theta$  and the value of  $v^2$  when  $P$  leaves the surface of the sphere. [7]

**END OF PAPER**