



**GCE**

**FURTHER MATHEMATICS**

**UNIT 6: FURTHER MECHANICS B**

**SAMPLE ASSESSMENT MATERIALS**

**(1 hour 45 minutes)**

### **ADDITIONAL MATERIALS**

In addition to this examination paper, you will need:

- a 12 page answer book;
- a Formula Booklet;
- a calculator.

### **INSTRUCTIONS TO CANDIDATES**

Use black ink or black ball-point pen.

Answer **all** questions.

Take  $g$  as  $9.8 \text{ ms}^{-2}$ .

Sufficient working must be shown to demonstrate the **mathematical** method employed.

Unless the degree of accuracy is stated in the question, answers should be rounded appropriately.

### **INFORMATION FOR CANDIDATES**

The number of marks is given in brackets at the end of each question or part-question. You are reminded of the necessity for good English and orderly presentation in your answers.

1. A ball of mass 0.4 kg is thrown vertically upwards from a point  $O$  with initial speed  $17 \text{ ms}^{-1}$ . When the ball is at a height of  $x$  m above  $O$  and its speed is  $v \text{ ms}^{-1}$ , the air resistance acting on the ball has magnitude  $0.01v^2 \text{ N}$ .

- (a) Show that, as the ball is ascending,  $v$  satisfies the differential equation

$$40v \frac{dv}{dx} = -(392 + v^2). \quad [3]$$

- (b) Find an expression for  $v$  in terms of  $x$ . [7]
- (c) Calculate, correct to two decimal places, the greatest height of the ball. [2]
- (d) State, with a reason, whether the speed of the ball when it returns to  $O$  is greater than  $17 \text{ ms}^{-1}$ , less than  $17 \text{ ms}^{-1}$  or equal to  $17 \text{ ms}^{-1}$ . [2]

2. (a) Prove that the centre of mass of a uniform solid cone of height  $h$  and base radius  $b$  is at a height of  $\frac{1}{4}h$  above its base. [4]

- (b) A uniform solid cone  $C_1$  has height 3 m and base radius 2 m. A smaller cone  $C_2$  of height 2 m and base radius 1 m is contained symmetrically inside  $C_1$ . The bases of  $C_1$  and  $C_2$  have a common centre and the axis of  $C_2$  is part of the axis of  $C_1$ . If  $C_2$  is removed from  $C_1$ , show that the centre of mass of the remaining solid is at a distance of  $\frac{11}{5}$  m from the vertex of  $C_1$ . [6]

- (c) The remaining solid is suspended from a string which is attached to a point on the outer curved surface at a distance of  $\frac{1}{3}\sqrt{13}$  m from the vertex of  $C_1$ .  
Given that the axis of symmetry is inclined at an angle of  $\alpha$  to the vertical, find  $\tan \alpha$ . [5]


3. A body, of mass 9 kg, is projected along a straight horizontal track with an initial speed of  $20 \text{ ms}^{-1}$ . At time  $t$  s the body experiences a resistance of magnitude  $(0.2 + 0.03v) \text{ N}$  where  $v \text{ ms}^{-1}$  is its speed.

- (a) Show that  $v$  satisfies the differential equation

$$900 \frac{dv}{dt} = -(20 + 3v). \quad [3]$$


- (b) Find an expression for  $t$  in terms of  $v$ . [5]
- (c) Calculate, to the nearest second, the time taken for the body to come to rest. [2]

4. The diagram shows a uniform lamina consisting of a rectangular section  $GPQE$  with a semi-circular section  $EFG$  of radius 4 cm. Quadrants  $APB$  and  $CQD$  each with radius 2 cm are removed. Dimensions in cm are as shown in the diagram.



- (a) Write down the distance of the centre of mass of the lamina  $ABCDEFG$  from  $AG$ . [1]
- (b) Determine the distance of the centre of mass of the lamina  $ABCDEFG$  from  $BC$ . [7]
- (c) The lamina  $ABCDEFG$  is suspended freely from the point  $E$  and hangs in equilibrium. Calculate the angle  $EG$  makes with the vertical. [3]

5. A particle  $A$ , of mass  $m$  kg, has position vector  $11\mathbf{i} + 6\mathbf{j}$  and a velocity  $2\mathbf{i} + 7\mathbf{j}$ . At the same moment, second particle  $B$ , of mass  $2m$  kg, has position vector  $7\mathbf{i} + 10\mathbf{j}$  and a velocity  $5\mathbf{i} + 4\mathbf{j}$ .
- (a) If the particles continue to move with these velocities, prove that the particles will collide. Given that the particles coalesce after collision, find the common velocity of the particles after collision. [9]
- (b) Determine the impulse exerted by  $A$  on  $B$ . [2]
- (c) Calculate the loss of kinetic energy caused by the collision. [2]
6. The diagram shows a playground ride consisting of a seat  $P$ , of mass 12 kg, attached to a vertical spring, which is fixed to a horizontal board. When the ride is at rest with nobody on it, the compression of the spring is 0.05 m.



The spring is of natural length 0.75 m and modulus of elasticity  $\lambda$ .

- (a) Find the value of  $\lambda$ . [2]

The seat  $P$  is now pushed vertically downwards a further 0.05 m and is then released from rest.

- (b) Show that  $P$  makes Simple Harmonic oscillations of period  $\frac{\pi}{7}$  and write down the amplitude of the motion. [5]
- (c) Find the maximum speed of  $P$ . [2]
- (d) Calculate the speed of  $P$  when it is at a distance 0.03 m from the equilibrium position. [3]
- (e) Find the distance of  $P$  from the equilibrium position 1.6 s after it is released. [3]
- (f) State one modelling assumption you have made about the seat and one modelling assumption you have made about the spring. [2]