

## Old Exam Questions – Old Course


**Moments**

(M1 Winter 2006)

5. A **non-uniform** rod  $AB$ , of mass  $7.5$  kg and length  $8$  m, rests horizontally in equilibrium on two smooth supports at  $C$  and  $D$ , where  $AC = 1.5$  m and  $AD = 5.0$  m. The reaction of the support at  $D$  on the rod is  $56.7$  N.
- (a) Calculate the distance of the centre of gravity of the rod from  $C$ . [4]
- (b) Determine the reaction of the support at  $C$  on the rod. [2]

(M1 Summer 2006)

6. The diagram shows a uniform straight rod  $AB$ , of length  $3.8$  m, resting horizontally in equilibrium on two smooth supports at  $C$  and  $D$  with an object of mass  $2.2$  kg freely suspended from point  $B$ .



The mass of the rod is  $4.4$  kg,  $AC = 0.4$  m and  $AD = 2.6$  m. Calculate the magnitudes of the reactions at  $C$  and  $D$ . [7]

(M1 Winter 2007)

7. The diagram shows a uniform plank  $XY$ , of mass  $40$  kg and length  $3$  m, resting on two supports at  $P$  and  $Q$ , where  $XP = 0.7$  m, and  $QY = 0.9$  m.




A boy  $A$ , of mass  $45$  kg, sits on the plank at the point  $P$  and a boy  $B$ , of mass  $70$  kg, sits on the plank at the end  $Y$ .

- (a) Modelling the boys as particles, calculate the magnitudes of the normal reactions of the supports on the plank. [6]
- (b) State what would happen if  $A$  jumps off the plank. Give a reason for your answer. [2]

(M1 Summer 2007)


7. The diagram shows a uniform rod  $AB$ , of length  $1.6$  m and mass  $8$  kg, held horizontally in equilibrium by means of two small smooth cylindrical pegs  $X$  and  $Y$ , such that  $AX = XY = 0.3$  m. A body of mass  $5$  kg is attached to the rod at point  $B$ .



Find the magnitude of each of the forces exerted on the rod by the pegs  $X$  and  $Y$ . [7]

(M1 Winter 2008)


7. The diagram shows a uniform rod  $AB$  of length  $3$  m and mass  $9$  kg, with a particle, of mass  $2$  kg, attached at  $A$ . The rod is resting horizontally in equilibrium on two smooth supports at points  $P$  and  $Q$  of the rod, where  $AP = 1.2$  m and  $AQ = 2.6$  m.



- (a) Calculate the reactions at  $P$  and  $Q$ . [7]
- (b) When an additional particle, of mass  $3$  kg, is attached to the point  $R$  of the rod, the rod is on the point of turning about  $P$ . Calculate the distance  $AR$ . [3]

(M1 Summer 2008)


8. A uniform rod  $AB$ , of length  $20$  cm and weight  $6$  N, is supported by two smooth supports at  $P$  and  $Q$ , one on each side of its centre  $C$ , with  $PC = CQ = 3$  cm, as shown in the diagram.



A body, of weight  $5$  N, is placed on the rod at a point which is  $x$  cm from the centre  $C$  of the rod. Find the greatest value of  $x$  if equilibrium is maintained. [5]

(M1 Winter 2009)

7. The diagram shows a uniform rod  $AB$ , of mass  $0.3$  kg and length  $2.0$  m, with three particles, of masses  $0.2$  kg,  $0.4$  kg and  $0.5$  kg, attached to the points  $A$ ,  $C$  and  $B$  respectively, where  $AC = 0.6$  m. When the rod is suspended by a string attached to the point  $X$  of the rod, it rests horizontally in equilibrium.



- (a) Calculate the tension in the string. [3]
- (b) Determine the distance  $AX$ . [4]

(M1 Summer 2009)


7. The diagram shows a **non-uniform** rod  $AB$ , of length  $6$  m and mass  $40$  kg, resting horizontally in equilibrium on two smooth supports at  $P$  and  $Q$ , which are respectively  $2.5$  m and  $5.5$  m from  $A$ . The point  $C$  is the position of the centre of mass of the rod and  $AC = x$  m. The forces exerted on the rod by the supports at  $P$  and  $Q$  are **equal** in magnitude.



- (a) Find the magnitude of each of the forces exerted on the rod by the supports at  $P$  and  $Q$ . [2]
- (b) Calculate the value of  $x$ . [4]

(M1 Winter 2010)

8. The diagram shows a wind chime consisting of a horizontal uniform rod  $AC$ , suspended in equilibrium by means of a light string attached to the mid-point  $X$  of the rod, together with three objects hanging from the points  $A$ ,  $B$  and  $C$  of the rod.




The length of the rod  $AC$  is  $20$  cm and the length of  $AB$  is  $8$  cm. The masses of the objects hanging from  $A$ ,  $B$ ,  $C$  are  $0.1$  kg,  $M$  kg,  $0.4$  kg respectively. The mass of the rod is  $0.5$  kg.

- (a) Find the value of  $M$ . [4]
- (b) Calculate the tension in the string. [3]

(M1 Summer 2010)


7. A uniform rod  $AB$  is suspended horizontally from the ceiling by means of two vertical light inextensible strings  $XA$  and  $YB$  of equal length.



The rod  $AB$  has mass 6 kg and length 1.4 m. A particle, of mass 10 kg, is attached to the rod at point  $C$ , where  $AC = 0.3$  m. Calculate the tension in **each** of the strings  $XA$  and  $YB$ . [7]

(M1 Winter 2011)


7. A uniform rod  $AB$ , of mass 3 kg, has length 2 m. A particle of mass 5 kg is attached to the end  $A$ , and a particle of mass 2 kg is attached to the end  $B$ . The diagram shows the rod resting horizontally in equilibrium on a smooth support at the point  $C$ , where  $AC = x$  m.



Calculate the magnitude of the reaction of the support at  $C$  and the value of  $x$ . [6]

(M1 Summer 2011)

8. The diagram shows a uniform rod  $AB$ , of mass 4 kg and length 1.6 m, with a particle, of mass 0.5 kg, attached at a point  $C$  of the rod, where  $AC = 0.5$  m. The rod is resting horizontally in equilibrium on two smooth supports at points  $X$  and  $Y$  of the rod, where  $AX = 0.6$  m and  $AY = 1.2$  m.



- (a) Calculate the reaction at  $X$  and the reaction at  $Y$ . [7]
- (b) When an additional particle of mass  $M$  kg is attached to the point  $C$ , the rod is on the point of turning about  $X$ . Calculate the value of  $M$ . [4]

(M1 Winter 2012)

7. The diagram shows a body, of mass 65 kg, attached to the end  $B$  of a uniform rigid rod  $AB$  of length 4 m. The mass of the rod is 35 kg. The rod is held horizontally in equilibrium by two smooth cylindrical pegs, one at  $A$  and another at  $C$ , where  $AC = 1.2$  m.



- (a) Write down the moment of the weight of the rod about the point  $A$ .  
State your units clearly. [2]
- (b) Find the forces exerted on the rod at  $A$  and  $C$ . [6]

(M1 Summer 2012)

8. A light uniform rod  $AB$  has length 1.4 m. A particle of mass 5 kg is attached to end  $A$ , and a particle of mass 2 kg is attached to end  $B$ . The rod rests horizontally in equilibrium on a smooth support at  $C$ .

- (a) Calculate the reaction of the support at  $C$ . [2]
- (b) Find the distance  $AC$ . [4]

(M1 Winter 2013)

7. A uniform beam  $AB$ , of length 6 m, rests in a horizontal position on two smooth supports at  $C$  and  $D$ , where  $AC = 1$  m and  $BD = 1.2$  m, as shown in the diagram.



- (a) When a vertical force of magnitude 1800 N is applied upwards to the beam at the end  $A$ , the beam is about to tilt about the support at  $D$ .  
Determine the weight of the beam. [5]
- (b) The vertical force is now removed so that the beam is resting in equilibrium on the two supports. Calculate the magnitude of the reaction of each of the supports at  $C$  and  $D$  on the beam. [5]

(M1 Summer 2013)

5. The diagram shows a uniform plank  $AB$  of mass  $12\text{ kg}$  and length  $2\text{ m}$ . The plank rests horizontally in equilibrium on two supports at  $C$  and at  $D$ , where  $AC = 0.8\text{ m}$  and  $AD = x\text{ m}$ .



- (a) The reaction of the support on the plank at  $D$  has magnitude  $84\text{ N}$ .
- Determine the reaction of the support on the plank at  $C$ .
  - Calculate the value of  $x$ . [7]
- (b) A rock of mass  $M\text{ kg}$  is placed at  $A$  so that the plank is on the point of tilting about  $C$ . Calculate the value of  $M$ . [3]


(M1 Winter 2014)

7. A uniform plank  $AB$ , of length  $4.8\text{ m}$  and mass  $M\text{ kg}$ , is resting on two smooth supports at points  $X$  and  $Y$ , such that  $AX = BY = 1.2\text{ m}$ .

- (a) A person of mass  $84\text{ kg}$  stands on the plank at a point which is  $0.8\text{ m}$  from  $B$ . The reaction of the support at  $X$  is of magnitude  $156.8\text{ N}$ . Find
- the value of  $M$ ,
  - the magnitude of the reaction of the support at  $Y$ . [6]
- (b) The person of mass  $84\text{ kg}$  walks along the plank towards  $A$ . At the instant that the plank starts to tilt about  $X$ , find
- the magnitude of the reaction of the support at  $X$ ,
  - the distance of the person from  $X$ . [5]

(M1 Summer 2014)


4. The diagram shows a uniform rod  $AB$ , of length  $1.8\text{ m}$  and mass  $3\text{ kg}$ , held in horizontal equilibrium by two small fixed cylinders  $C$  and  $D$ . An object of mass  $12\text{ kg}$  rests on the rod at  $B$ . The length  $AC$  is  $0.3\text{ m}$  and  $CD$ , the distance between the cylinders, is  $0.4\text{ m}$ . The force exerted on the rod by each of the cylinders is vertical.



Find the magnitude of each of the forces exerted on the rod by the cylinders. [7]

(M1 Summer 2015)

5. The diagram shows a plank  $AB$ , of mass  $15\text{ kg}$  and length  $2.8\text{ m}$ , being held in equilibrium with  $AB$  horizontal by means of two vertical ropes, one attached to the end  $A$  and the other attached to the end  $B$ . A man of mass  $80\text{ kg}$  stands on the plank at point  $C$ , where  $AC = 0.9\text{ m}$ .



- (a) Modelling the plank as a uniform rod, find the tensions in the ropes attached to the end  $A$  and the end  $B$  of the plank. [7]
- (b) The plank is now modelled as a **non-uniform** rod. Given that the tension in the rope attached to  $A$  is  $1.5$  times the tension in the rope attached to  $B$ , determine the distance of the centre of mass of the plank from  $A$ . [5]

(M1 Summer 2016)

6. A uniform rod  $AB$  is of mass  $8\text{ kg}$  and length  $6\text{ m}$ . It is suspended horizontally in equilibrium by means of two vertical light strings attached to the rod  $AB$  at point  $C$  and point  $D$  on the rod, where  $AC = 1.6\text{ m}$  and  $AD = 4.8\text{ m}$ . Calculate the tension in the string at  $C$  and the tension in the string at  $D$ . [7]

(M1 Summer 2017)

3.



The diagram shows a uniform plank  $AB$ , of mass  $20\text{ kg}$  and length  $2.4\text{ m}$ , supported in horizontal equilibrium by two pivots, one at  $C$  and one at  $D$ . The distance  $AC$  and the distance  $DB$  are both  $0.5\text{ m}$ . A person of mass  $40\text{ kg}$  stands at a point which is  $0.6\text{ m}$  from  $B$ .

- (a) Calculate the magnitudes of the reaction at  $C$  and the reaction at  $D$ . [7]
- (b) The person starts to walk towards  $A$ . Determine the greatest distance of the person from  $B$  if equilibrium is to be maintained. [3]

(M1 Summer 2018)

3. A uniform rod  $AB$  is resting in horizontal equilibrium on two smooth supports at  $P$  and  $Q$ . The length of the rod is  $2\text{ m}$  and its mass is  $24\text{ kg}$ . Supports  $P$  and  $Q$  are such that  $AP = 0.3\text{ m}$  and  $PQ = 1.1\text{ m}$ . A person of mass  $36\text{ kg}$  stands on the rod at the end  $A$ . Find the magnitude of the reactions at  $P$  and at  $Q$ . [7]