



**GCE AS/A level**

0976/01

**MATHEMATICS C4**  
**Pure Mathematics**

A.M. THURSDAY, 14 June 2012

1½ hours

#### **ADDITIONAL MATERIALS**

In addition to this examination paper, you will need:

- a 12 page answer book;
- a Formula Booklet;
- a calculator.

#### **INSTRUCTIONS TO CANDIDATES**

Use black ink or black ball-point pen.

Answer **all** questions.

Sufficient working must be shown to demonstrate the **mathematical** method employed.

#### **INFORMATION FOR CANDIDATES**

The number of marks is given in brackets at the end of each question or part-question.

You are reminded of the necessity for good English and orderly presentation in your answers.

1. The function  $f$  is defined by

$$f(x) = \frac{11 + x - x^2}{(x + 1)(x - 2)^2}.$$

(a) Express  $f(x)$  in terms of partial fractions. [4]

(b) Use your result to part (a) to find the value of  $f'(0)$ . [3]

2. Find the equation of the tangent to the curve

$$y^3 - 4x^2 - 3xy + 25 = 0$$

at the point  $(2, -3)$ . [4]

3. (a) Find all values of  $\theta$  in the range  $0^\circ \leq \theta \leq 360^\circ$  satisfying

$$4 \cos 2\theta = 1 - 2 \sin \theta. \quad [6]$$

(b) (i) Express  $8 \sin x + 15 \cos x$  in the form  $R \sin(x + \alpha)$ , where  $R$  and  $\alpha$  are constants with  $R > 0$  and  $0^\circ < \alpha < 90^\circ$ .

(ii) Find all values of  $x$  in the range  $0^\circ \leq x \leq 360^\circ$  satisfying

$$8 \sin x + 15 \cos x = 11.$$

(iii) Find the greatest possible value for  $k$  so that

$$8 \sin x + 15 \cos x = k$$

has solutions. Give a reason for your answer. [7]

4. The region  $R$  is bounded by the curve  $y = \sqrt{x} + \frac{5}{\sqrt{x}}$ , the  $x$ -axis and the lines  $x = 3$ ,  $x = 4$ .

Find the volume generated when  $R$  is rotated through four right-angles about the  $x$ -axis. Give your answer correct to the nearest integer. [5]

5. Expand  $\left(1 + \frac{x}{3}\right)^{-\frac{1}{2}}$  in ascending powers of  $x$  up to and including the term in  $x^2$ .

State the range of values of  $x$  for which your expansion is valid.

Hence, by writing  $x = \frac{1}{5}$  in your expansion, find an approximate value for  $\sqrt{15}$  in the form  $\frac{a}{b}$ , where  $a$  and  $b$  are integers whose values are to be found. [5]

6. The parametric equations of the curve  $C$  are  $x = t^2$ ,  $y = 2t$ .

(a) Show that the normal to  $C$  at the point  $P$  with parameter  $p$  has equation

$$y + px = p^3 + 2p. \quad [5]$$

(b) The normal to  $C$  at the point  $P$  intersects  $C$  again at the point with parameter 3.

(i) Show that  $p^3 - 7p - 6 = 0$ .

(ii) Hence show that  $P$  can be one of two points. Find the coordinates of each of these two points. [6]

7. (a) Find  $\int xe^{-2x} dx$ . [4]

(b) Use the substitution  $u = 1 + 3\ln x$  to evaluate

$$\int_1^e \frac{1}{x(1 + 3\ln x)} dx.$$

Give your answer correct to four decimal places. [4]

8. Water is leaking from a hole at the bottom of a large tank. The volume of the water in the tank at time  $t$  hours is  $V \text{ m}^3$ . The rate of decrease of  $V$  is directly proportional to  $V^3$ .

(a) Write down a differential equation satisfied by  $V$ . [1]

(b) Given that  $V = 60$  when  $t = 0$ , show that

$$V^2 = \frac{3600}{at + 1},$$

where  $a$  is a constant. [4]

(c) When  $t = 2$ , the volume of the water in the tank is  $50 \text{ m}^3$ . Find the value of  $t$  when the volume of the water in the tank is  $27 \text{ m}^3$ . Give your answer correct to one decimal place. [4]

## TURN OVER

9. The position vectors of the points  $A$  and  $B$  are given by

$$\mathbf{a} = 4\mathbf{i} + \mathbf{j} - 6\mathbf{k},$$

$$\mathbf{b} = 6\mathbf{i} + 2\mathbf{j} - 4\mathbf{k},$$

respectively.

- (a) Determine whether or not the vectors  $\mathbf{a}$  and  $\mathbf{b}$  are perpendicular, giving a reason for your answer. [2]

- (b) (i) Write down the vector  $\mathbf{AB}$ .  
 (ii) Find the vector equation of the line  $AB$ . [3]

- (c) The vector equation of the line  $L$  is given by

$$\mathbf{r} = 2\mathbf{i} + 6\mathbf{j} + p\mathbf{k} + \mu(-2\mathbf{i} + \mathbf{j} + 3\mathbf{k}),$$

where  $p$  is a constant.

Given that the lines  $AB$  and  $L$  intersect, find the value of  $p$ . [5]

10. Complete the following proof by contradiction to show that  $\sqrt{5}$  is irrational.

*Assume that  $\sqrt{5}$  is rational. Then  $\sqrt{5}$  may be written in the form  $\frac{a}{b}$ , where  $a, b$  are integers having no common factors.*

$$\therefore a^2 = 5b^2.$$

$\therefore a^2$  has a factor 5.

$\therefore a$  has a factor 5 so that  $a = 5k$ , where  $k$  is an integer. [3]