



**GCE AS/A level**

0979/01



S16-0979-01

**MATHEMATICS – FP3**  
**Further Pure Mathematics**

A.M. WEDNESDAY, 29 June 2016

1 hour 30 minutes

**ADDITIONAL MATERIALS**

In addition to this examination paper, you will need:

- a 12 page answer book;
- a Formula Booklet;
- a calculator.

**INSTRUCTIONS TO CANDIDATES**

Use black ink or black ball-point pen.

Answer **all** questions.

Sufficient working must be shown to demonstrate the **mathematical** method employed.

**INFORMATION FOR CANDIDATES**

The number of marks is given in brackets at the end of each question or part-question.

You are reminded of the necessity for good English and orderly presentation in your answers.

1. The curve  $C$  has polar equation

$$r = 1 + 2 \tan \theta, \quad 0 \leq \theta \leq \frac{\pi}{4}.$$

Show that there is no point on  $C$  at which the tangent is perpendicular to the initial line. [7]

2. The function  $f$  is defined by  $f(x) = \cos x + \cosh x$ .

(a) Show that  $f^{(4)}(x) = f(x)$ , where  $f^{(4)}(x)$  denotes the fourth derivative of  $f(x)$ . [2]

(b) (i) Show that the Maclaurin series of  $f(x)$  contains only terms of the form  $x^{4n}$ , where  $n$  is a non-negative integer.

(ii) Determine the first three non-zero terms of this Maclaurin series. [3]

(c) (i) Hence find an approximate value for the positive root of the equation

$$12(\cos x + \cosh x) - x^4 = 36.$$

Give your answer correct to three significant figures.

(ii) Show that this approximation is the value of the root correct to three significant figures. [5]

3. Using the substitution  $t = \tan\left(\frac{x}{2}\right)$ , evaluate the integral

$$\int_0^{\frac{\pi}{2}} \frac{dx}{3 + 5\cos x},$$

giving your answer in the form  $\ln(3^a)$ , where  $a$  is a rational number to be determined. [8]

4. The function  $f$  is defined on the domain  $[0, \infty)$  by

$$f(\theta) = \cosh 2\theta - 8\cosh \theta.$$

Consider the equation  $f(\theta) = k$ , where  $k$  is a constant.

(a) Show that the equation has no real roots if  $k < -9$ . [4]

(b) Solve the equation when  $k = -8$ , giving your answers correct to two decimal places. [3]

(c) Determine

(i) the value of  $k$  for which the equation has a repeated root,

(ii) the set of values of  $k$  for which the equation has exactly one real root. [5]

5. The curve  $C$  has equation  $y = \ln(1 + \cos x)$ .

(a) Show that

$$1 + \left(\frac{dy}{dx}\right)^2 = \frac{2}{1 + \cos x} . \quad [4]$$

(b) Find the length of the arc joining the points  $(0, \ln 2)$  and  $\left(\frac{\pi}{2}, 0\right)$  on  $C$ .

Give your answer in the form  $\ln(a + b\sqrt{2})$ , where  $a, b$  are positive integers. [6]

6. The equation

$$x^5 + \sinh x = 3$$

has a root  $\alpha$  close to 1.

(a) It is suggested that iterative sequences based on the following rearrangements of the equation could be used to find the value of  $\alpha$ .

I.  $x = (3 - \sinh x)^{\frac{1}{5}}$

II.  $x = \sinh^{-1}(3 - x^5)$

(i) By evaluating appropriate derivatives, show that one of these sequences is convergent and the other is divergent.

(ii) Taking  $x_0 = 1$ , use the convergent sequence to find the value of  $\alpha$  correct to three decimal places. [12]

(b) Use the Newton-Raphson method to find the value of  $\alpha$  correct to six decimal places. [6]

7. The integral  $I_n$  is given, for  $n \geq 0$ , by

$$I_n = \int_0^{\pi} x^n \sin 2x \, dx.$$

(a) Show that, for  $n \geq 2$ ,

$$I_n = -\frac{\pi^n}{2} - \frac{n(n-1)}{4} I_{n-2} . \quad [6]$$

(b) Evaluate  $I_4$ , giving your answer correct to the nearest integer. [4]

**END OF PAPER**