



MARKING SCHEME

**LEVEL 2 CERTIFICATE IN ADDITIONAL
MATHEMATICS**

SUMMER 2015

INTRODUCTION

The marking schemes which follow were those used by WJEC for the Summer 2015 examination in LEVEL 2 CERTIFICATE IN ADDITIONAL MATHEMATICS. They were finalised after detailed discussion at examiners' conferences by all the examiners involved in the assessment. The conferences were held shortly after the papers were taken so that reference could be made to the full range of candidates' responses, with photocopied scripts forming the basis of discussion. The aim of the conferences was to ensure that the marking schemes were interpreted and applied in the same way by all examiners.

It is hoped that this information will be of assistance to centres but it is recognised at the same time that, without the benefit of participation in the examiners' conferences, teachers may have different views on certain matters of detail or interpretation.

WJEC regrets that it cannot enter into any discussion or correspondence about these marking schemes.

LEVEL 2 CERTIFICATE IN ADDITIONAL MATHEMATICS
Mark Scheme - Summer 2015

	Additional Mathematics Summer 2015		Final
1	$(2x - 5)(3x + 2)$ $5/2$ and $-2/3$	B2 B2 4	B1 $(2x \dots 5)(3x \dots 2)$ or $(2x \dots 2)(3x \dots 5)$ Ignore sight of “=0” Must be from factorising, do not accept use of quadratic formula followed by ‘factorising’. MUST FT for their factors FT for their factors equivalent difficulty not leading to whole number solutions. B1 for each answer Allow -0.66 or -0.7 as a solution provided $3x = -2$ is seen in working
2	(a) $(x + 7)^2$ ($\pm \dots$) (Minimum value at $x =$) -7 (b) -40	M1 A1 B1 3	Ignore ‘their ($\pm \dots$)’ or ‘=0’ Do not accept method $dy/dx = 2x + 14$ Unsupported -7 is M0, A0 CAO Must be given in (b), do not accept shown in (a). Do not accept reversed answers of ‘-40’ in (a) and ‘-7’ in (b), maximum mark would be possible M1 in (a) However, if no marks in (a), but a full statement, e.g. ‘minimum value -40 when $x = -7$ ’ is given in (b), then award B1 in (b) and M1, A0 in (a)
3	(a) $40x^7 - 6$ (+0) (b) $-8x^{-9}$ (or $-8/x^9$) (c) $2/5x^{-3/5}$ or equivalent	B3 B1 B1 5	B1 for $40x^7$ (not $5 \times 8x^7$), B1 for -6 and B1 for +0 (or blank) provided at least one other mark awarded. CAO. Index needs to be simplified. CAO. Index needs to be simplified. <u>ISW once simplified to stages shown in (b) and (c)</u> Penalise including ‘+c’ -1 only
4	(a) $(DE^2 =) (6 - 4)^2 + (22 - 14)^2 (=10^2 + 8^2)$ $DE = \sqrt{164}$ $= 2\sqrt{41}$ (b) Gradient DE $(22 - 14) / (6 - 4)$ $= 8/10 (= 4/5)$ Gradient perpendicular $-10/8 (= -5/4)$ $(6 + 4)/2, (22 + 14)/2$ Mid point DE $(1, 18)$ or equivalent Use of $y = mx + c$ or $y - y_1 = m(x - x_1)$ $y = -10x/8 + 19\frac{1}{4}$ or $y - 18 = -10/8 (x - 1)$ $10x + 8y - 154 = 0$ OR $5x + 4y - 77 = 0$ OR $-10x - 8y + 154 = 0$ OR $-5x - 4y + 77 = 0$	M1 A1 B1 M1 A1 B1 M1 A1 M1 A1 A1 11	Or $(-4 - 6)^2 + (14 - 22)^2$. Allow 1 slip in sign of substitution CAO FT ‘their DE’ of equivalent difficulty expressed correctly, e.g. $\sqrt{18} = 3\sqrt{2}$, needs to be in the form $a\sqrt{b}$ where $a \neq 1$ and $b \neq 1$ or simpler Sight of $2\sqrt{41}$ implies previous $\sqrt{164}$ Or $(14 - 22)/(-4 - 6)$ CAO. Mark final answer and then FT FT $-1/\text{grad DE}$ Accept $(1, \dots)$ or $(\dots, 18)$ CAO Method to find the equation using mid-point and perpendicular gradient (not $8/10$) FT their mid-point (not D or E) & their perpendicular gradient, or FT substitution of their midpoint with their perpendicular gradient in $y = mx + c$ (towards finding c) <i>If no working for finding gradient is seen, then ‘their ‘spurious’ incorrect perpendicular gradient’ must be negative</i> FT for correct unsimplified form, not written in quotient form CAO. Form $ax + by + c = 0$ or a rearrange of this provided it is ‘=0’

	Additional Mathematics Summer 2015		Final
5	<p>Sight of $x-4$ and $x+1$</p> <p>$2(x(x-4) + x(x+1) + (x-4)(x+1)) = 124$</p> <p>$2x^2 - 8x + 2x^2 + 2x + 2x^2 - 8x + 2x - 8 = 124$ OR $x^2 - 4x + x^2 + x + x^2 - 4x + x - 4 = 62$</p> <p>$6x^2 - 12x - 8 = 124$ OR $3x^2 - 6x - 4 = 62$</p> <p>Shows or sight of $x^2 - 2x - 22$ or $x^2 - 2x - 22 = 0$</p> <p>$(x - 1)^2 = 22 + 1$ OR $x = \frac{2 \pm \sqrt{92}}{2}$</p> <p>$x = 1 + \sqrt{23}$</p> <p>QWC2:</p> <ul style="list-style-type: none"> Candidates will be expected to present work clearly, with words explaining process or steps <p>AND</p> <ul style="list-style-type: none"> make few if any mistakes in mathematical form, spelling, punctuation and grammar in their answer <p>QWC1: Candidates will be expected to</p> <ul style="list-style-type: none"> present work clearly, with words explaining process or steps <p>OR</p> <ul style="list-style-type: none"> make few if any mistakes in mathematical form, spelling, punctuation and grammar in their final answer 	<p>B1</p> <p>M2</p> <p>m1</p> <p>A1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>QWC 2</p> <p>10</p>	<p><i>FT 1 error in 1 side length, provided it is in the form 'x ± a' and not a single term expression, for M marks only</i></p> <p><i>Some work may be shown in stages, but implies the following M and m marks, but need to show summation</i></p> <p>Correct expression for the total surface area Accept '×2' omitted if equated to 62 M1 for '×2' aspect omitted, or M1 for '×2' included with any 2 (of 3) area expressions correct, or M1 for '×2' included with any 4 (of 6) areas correct, or M1 if it could be correct but brackets missing and not implied in later working (if implied in later working allow M2)</p> <p>Equate to 124 with '×2' treated correctly, or equate to 62 if appropriate, with at least 2 of the 3 (or 4 of the 6) brackets expanded correctly; depends on M1</p> <p>Depends on M2, m1 for correct manipulation and collection of terms</p> <p>Depends on M2, m1. Requirement to show as asked in the question</p> <p>Independent of other marks Must be correct to this stage of working FT for their quadratic equation of equivalent difficulty</p> <p>CAO. Do not accept $1 \pm \sqrt{23}$</p> <p>QWC2 Presents relevant material in a coherent and logical manner, using acceptable mathematical form, and with few if any errors in spelling, punctuation and grammar.</p> <p>QWC1 Presents relevant material in a coherent and logical manner but with some errors in use of mathematical form, spelling, punctuation or grammar OR evident weaknesses in organisation of material but using acceptable mathematical form, with few if any errors in spelling, punctuation and grammar.</p> <p>QWC0 Evident weaknesses in organisation of material, and errors in use of mathematical form, spelling, punctuation or grammar.</p>

	Additional Mathematics Summer 2015		Final
6	$2x+1 = x^2 + 6x - 5$ $x^2 + 4x - 6 = 0$ $x = \{-4 \pm \sqrt{(4^2 - 4 \times 1 \times -6)}\}/2$ $x = \{-4 \pm \sqrt{40}\}/2$ $x = 1.16$ and $x = -5.16$ $x = 1.16$ with $y = 3.32$ or $y = 3.31$ and $x = -5.16$ with $y = -9.32$ or $y = -9.33$	M1 A1 m1 A1 A1 A1 6	<p>Must be equated to zero. ‘=0’ may be implied in further work to solve, if no further work and not ‘=0’ then A0</p> <p>FT provided their quadratic does not factorise and equivalent level of difficulty Use of correct quadratic formula, allow 1 slip in substitution (not a slip with the formula) If completing the square used award m1 for sight of $(x + 2)^2 \pm \dots$</p> <p>FT provided M1,m1 previously awarded using their values of x in $2x + 1$ or equivalent to find y-values to 2 d.p.</p> <p><i>Alternative using $x = (y - 1)/2$</i></p> <p>M1 $y = \frac{(y-1)^2 + 6(y-1) - 5}{+1/4}$ or $y = y^2/4 + 5y/2 - 8$ A1 $y^2 + 6y - 31 = 0$ or equivalent (equate to zero) m1 $y = \{-6 \pm \sqrt{(6^2 - 4 \times 1 \times -31)}\}/2$ or equivalent ($\times 1/4$) Allow 1 slip in substitution A1 $y = (-6 \pm \sqrt{160})/2$ or equivalent A1 $y = 3.32$ and $y = -9.32$ A1 $x = 1.16, y = 3.32$ and $x = -5.16, y = -9.32$ FT to final A1, provided M1,m1 previously awarded using their values of y in $(y - 1)/2$ or equivalent to find x-values to 2 d.p.</p>
7	(a) $3(-2)^3 - 2(-2)^2 + 5(-2) - 1$ (= -24-8-10-1) -43 (b)(i) Substitute $x = 2$ Showing $f(2) = 0$ (ii) $(x-2)(x^2 + bx + c)$ or intention to divide by $(x-2)$ with x^2 shown $((x-2)) (x^2 + 10x + 21)$ $((x-2)) (x + 3)(x + 7)$	M1 A1 M1 A1 M1 A2 A1 8	<p>Or division method giving $3x^2 - 8x \dots$</p> <p>Or division method giving $x^2 + 10x \dots$ Accept sight of substitution with ‘=0’ shown</p> <p>If any values are inserted at least 1 needs to be correct, appropriate sight of $+10x$ or $+21$ implies M1 (and A1 to follow)</p> <p>A1 for $(+10x$ or $(+21$ Or use of factor theorem A1 $(x+3)$, A1 $(x+7)$</p> <p>CAO, but ignore sight of “=0”, ISW</p>
8	(a) $180x^8$ (b) $a = 1$ $c = 5$ $b = 3$	B2 B1 B1 B1 5	<p>B1 for sight of $20x^9$. FT to 2nd B1 from $dy/dx = kx^n$ Ignore incorrect notation</p> <p>FT $b = 8 -$ ‘their c’ or $b = 4 -$ ‘their a’</p> <p><i>Accept sight of correct answers from ‘uncorrected’ working</i> Do not accept embedded answers, candidates need to identify values for a, b and c, not accept as left in working without clearly stating.</p>

	Additional Mathematics Summer 2015		Final
9	(a) $21x^{7/7} - 3x^3/3 - x^{-1}/-1 + 6x + c$ (constant) (b) $6x^3/3 + 4x^2/2$ $[6x^3/3 + 4x^2/2]^5_2$ $= (2 \times 5^3 + 2 \times 5^2) - (2 \times 2^3 + 2 \times 2^2)$ $= 276$	B4 B1 B2 M1 A1 A1 10	B1 for each term. Accept unsimplified $-x^{-1}/-1$. ISW Awarded if at least B1 for integration Mark final answer, then FT. B1 for sight of $6x^3/3$ or $4x^2/2$ FT their integration, not original. Intention to use 5, 2 and subtract FT for correct use of limits Accept unsimplified fractions included CAO, not FT. Do not accept ' $276 + c$ ' <i>Answer only, no working shown M0 A0 A0</i>
10	(a) $(\frac{1}{2})^2 + (\sqrt{3}/2)^2$ $\frac{1}{4} + \frac{3}{4} (=1)$ (b) $5 \times 1 + 2 \times \sqrt{3}/2 + \sqrt{3}$ $5 + 2\sqrt{3}$	M1 A1 M2 A1 5	Use of $\frac{1}{2}$ and $\sqrt{3}/2$ appropriately in either order M1 for any 2 terms correct M1 only if first line of working is $5 + \sqrt{3} + \sqrt{3}$, but allow FT for possible A1 CAO. Mark final answer
11	3D visualisation with height 12cm AND use of appropriate methods Side length of base 8cm or $\frac{1}{2}$ side 4cm $\text{diagonal}^2 = 8^2 + 8^2$ OR $(\frac{1}{2} \text{diagonal})^2 = 4^2 + 4^2$ diagonal is $\sqrt{128}$ or $8\sqrt{2}$ (cm) OR $\frac{1}{2}$ diagonal is $\frac{1}{2}\sqrt{128}$ or $4\sqrt{2}$ tan 'required angle' = $(12 / \frac{1}{2}\text{diagonal})$ or full alternative method $64.7(6\dots^\circ)$ or 64.8°	S1 B1 M1 A1 M1 A2 7	e.g. 3D sketch with sight of height 12cm, Pythagoras' Theorem and tan ratio, or full alternative trig method following Pythagoras' Theorem, or use of base area to find the side length followed by full alternative trig method Sight of 8cm or 4cm as appropriate FT their side length, provided $\neq 64$, throughout other than the final A mark. (diagonal= $11.3137\dots$ cm, $\frac{1}{2}$ diagonal = $5.65685\dots$ cm rounded or truncated) FT their derived $\frac{1}{2}$ diagonal for M1 only, not 8 or 4(cm) Full alternative method: finding perpendicular height of a sloping face then the slant edge CAO. Accept 65° from correct working A1 for $\tan^{-1}(12/\frac{1}{2}\sqrt{128})$, or for an answer from premature approximation
12	$y + \delta y = (x + \delta x)^2 - 3(x + \delta x)$ Intention to subtract $(y =) x^2 - 3x$ to find δy $\delta y = 2x\delta x + (\delta x)^2 - 3\delta x$ Dividing by δx and $(\lim) \delta x \rightarrow 0$ $dy/dx = \lim_{\delta x \rightarrow 0} \delta y/\delta x = 2x - 3$	B1 M1 A1 M1 A1 5	Or alternative notation. Allow if final bracket omitted Accept δx^2 as meaning $(\delta x)^2$ FT equivalent level of difficulty CAO. Must follow from correct working <i>Use of dy/dx throughout or incorrect notation then possible maximum is only 4 marks, final A0</i>
13	When $x = 3$, finding $y = -6$ $dy/dx = 4x - 8$ when $x = 3$ gradient is 4 Use of $y - y_1 = m(x - x_1)$ or $y = mx + c$ $y - -6 = 4(x - 3)$ or $-6 = 4 \times 3 + c, c = -18$ $4x - y - 18 = 0$	B1 M1 A1 M1 A1 A1 6	Method to form equation FT their y value, but not $y=3$ and their derived gradient CAO. Must be in this form, accept equivalents written as 3 terms with whole number coefficients with ' $=0$ ' or ' $0=$ '

	Additional Mathematics Summer 2015		Final
14	$(dy/dx) 6x^2 - 24$ $dy/dx = 0$ or $6x^2 - 24 = 0$ $x = 2$ and $y = -19$ $x = -2$ and $y = 45$ $d^2y/dx^2 = 12x$ At $(2, -19)$ $d^2y/dx^2 > 0$, point is a minimum At $(-2, 45)$: $d^2y/dx^2 < 0$, point is a maximum	B1 M1 A1 A1 M1 A1 A1 7	FT their dy/dx form $ax^2 + b$ throughout <i>Answer only, no working shown M0 A0 A0</i> <i>Method for determining min or max MUST be shown, final answer only is M0 here, then A0,A0</i> Or first derivative test, interpretation of first derivative test. Or alternative. FT for their x value FT for their other x value provided this does not have the same interpretation as the first x value <i>SC1 for correct FT from $d^2y/dx^2 = ax, a > 0$</i> <i>Do not accept trial & improvement methods unless both stationary points are found correctly and confirmed as stated in the mark scheme</i>
15	(a) General cos curve intersecting x-axis only at $(90^\circ, 0)$ and $(270^\circ, 0)$ Correct curve with 5 and -5 on y-axis (b) 90° and 270° only	M1 A1 B1 3	Allow general shape as the joining of key values, but straight rather than clearly curving towards a turn at 0° and 360° in particular Must show a clear curve, not a straight a 0° and 360° in particular This values need to be selected, not amongst others unless unambiguous indicated as the response.
16	144	B1 1	<i>No marks if no working. Must see 12^2 or $(2\sqrt{3})^4 = 16 \times 9$</i>
17	(a) $(20)x^{8/8}/x^{2/3}$ or equivalent first stage of work evaluated correctly with simplification of indices $20x^{1/3}$ or $20\sqrt[3]{x}$ (b) Correctly extracting a factor of $x^{1/4}$ (to give correct numerator) OR correct alternative method with one correct step towards simplification $2 + x^{1/2}$ or $2 + \sqrt{x}$	B1 B1 M1 A1 4	 CAO. Mark final answer For an alternative method, need sight of the two terms and $2 + \dots$ or $\dots + x^{2/4}$ for M1 CAO accepting $2 + x^{2/4}$. Mark final answer



WJEC
245 Western Avenue
Cardiff CF5 2YX
Tel No 029 2026 5000
Fax 029 2057 5994
E-mail: exams@wjec.co.uk
website: www.wjec.co.uk